

1 **Identification and Utility of Native Fish Conservation Areas**
2 **in the Upper Snake River Basin, USA**

3
4
5 RICHARD N. WILLIAMS*

6 *Department of Biology, The College of Idaho, 2112 Cleveland Blvd, Caldwell, Idaho 83605, USA*

7
8 DANIEL C. DAUWALTER

9 *Trout Unlimited, 910 Main Street, Suite 342, Boise, Idaho 83702, USA*

10
11 RUSSELL F. THUROW

12 *USDA Forest Service-Rocky Mountain Research Station, 322 East Front Street- Suite 401, Boise,*
13 *Idaho 83702, USA*

14
15 DAVID P. PHILIPP

16 *Fisheries Conservation Foundation, 302 E. Green Street #2102, Champaign, Illinois 61825, USA*

17
18 JACK E. WILLIAMS

19 *Trout Unlimited, 4393 Pioneer Road, Medford, Oregon 97501, USA*

20
21 CHRIS A. WALSER

22 *Department of Biology, The College of Idaho, 2112 Cleveland Blvd, Caldwell, Idaho 83605, USA*

23
24 *Corresponding author: troutdna@gmail.com

27 *Abstract.*--Native Fish Conservation Areas (NFCAs) are watersheds where management
28 emphasizes proactive conservation and restoration for long-term persistence of native fish
29 assemblages while allowing for compatible uses. NFCAs are intended to complement traditional
30 fisheries management approaches that are often reactive to population stressors and focused
31 on single species conservation efforts rather than complete assemblages. We identified
32 potential NFCAs in the Upper Snake River basin above Hells Canyon Dam using a process that
33 ranked all subwatersheds (Hydrologic Unit Code 12) and used empirical data on distribution,
34 abundance, and genetics for three native trout species (Bull Trout *Salvelinus confluentus*;
35 Columbia River Redband Trout *Oncorhynchus mykiss*; and Yellowstone Cutthroat Trout *O.*
36 *clarkii*, including the fine-spotted form) and both known occurrences and modeled potential
37 distributions of native non-game fishes. Rankings also incorporated drainage network
38 connectivity and land protection status (e.g., national park, wilderness). Clusters of high-ranking
39 subwatersheds were identified as potential NFCAs that were then classified according to the
40 presence of non-game fishes identified as Species of Greatest Conservation Need in state
41 wildlife action plans. Pacific Creek and Goose Creek watershed ranked high in the upper basin
42 (above Shoshone Falls), and Little Jacks Creek and Squaw Creek ranked high in the lower basin.
43 We then contrasted characteristics of a select few potential NFCAs; discuss the practical
44 implementation and benefits of NFCAs for both fishes and other aquatic species in the Upper
45 Snake River basin; examine how the NFCA approach could enhance existing conservation
46 partnerships; and discuss how designating select watersheds as NFCAs can create higher public
47 awareness of the value of native fishes and other aquatic species and their habitats.

48
49

Introduction

50 Despite decades of allocating substantial resources to conserve freshwater ecosystems,
51 North American freshwater fishes continue to decline at a much faster rate than their
52 terrestrial counterparts (Master et al. 2000; Jelks et al. 2008). Williams (2019; Chapter 1)
53 discusses how current conservation approaches, such as the National Wildlife Refuge system,
54 have been only moderately successful in protecting riverine ecosystems. Because rivers are
55 linear in nature, approaches based on terrestrial features and land ownership often fail to
56 consider watershed boundaries fundamental to aquatic conservation (Saunders et al. 2002;
57 Roux et al. 2008).

58 Williams et al. (2011) proposed the concept of Native Fish Conservation Areas (NFCAs) to
59 establish entire watersheds cooperatively managed for native fish communities. As a
60 complement to existing conservation approaches (e.g., headwater isolation; Novinger and
61 Rahel 2003), NFCAs emphasize intact and persistent native fish communities and healthy and
62 resilient ecosystems while simultaneously striving to support compatible commercial and
63 recreational uses. Dauwalter et al. (2011) explored the NFCA concept and its application in the
64 Upper Colorado River Basin in Wyoming through a process that combined known and modeled
65 species distributions, spatial prioritization analysis, and stakeholder discussions. Others have
66 used the NFCA concept as an organizing framework for broad-scale native fish conservation
67 initiatives and associated funding programs (Birdsong et al. 2015; Birdsong et al. 2019, Chapter
68 X).

69 In this paper, we further explore the utility of the NFCA concept by identifying potential
70 NFCAs and their benefits in the Upper Snake River basin upstream from Hells Canyon Dam. As
71 described in detail below, we used known distributions and abundance of native trout species,
72 known and modeled occurrences of non-game fishes, drainage network connectivity
73 interrupted by large dams, and land protection status (e.g., national parks, wilderness, wild and
74 scenic river) to rank all subwatersheds based on their conservation value and identify potential
75 NFCAs. We then summarize information by potential NFCAs, compare implementation of
76 NFCAs in example watersheds, and discuss the utility of the NFCAs in the Upper Snake River
77 basin.

78

79

Upper Snake River Basin

80 The Snake River flows through a large basin draining portions of Wyoming, Idaho, Utah,
81 Nevada, and Oregon. From its headwaters in Yellowstone National Park, Wyoming, the river
82 flows southwest, then west, and then northwest before cascading north through Hells Canyon
83 along the Idaho-Oregon border (Figure 1).

84 The Upper Snake basin lies above the Hells Canyon Complex of three dams (Brownlee,
85 Oxbow, and Hells Canyon dams), that is the present upstream limit to anadromous salmon and
86 steelhead migrations into the Snake River in Idaho. The Upper Snake basin is naturally divided
87 by Shoshone Falls, a 65 m waterfall near Twin Falls, Idaho that effectively separates the Snake
88 River basin into upper and lower basins and is a complete barrier to fish migration (Figure 1;
89 Behnke and Cushing 2005). The falls, along with a unique geologic history and past connections
90 with Pleistocene Lake Bonneville, have resulted in a unique history of species colonization that
91 has strongly influenced the biogeography of fishes in the Snake River basin (Smith 1978;
92 Campbell et al. 2011). The lower basin (Shoshone Falls to Hells Canyon Dam) supported 25
93 native fish species of which five are extirpated (Table 1), compared to the upper basin
94 (Yellowstone Park headwaters to Shoshone Falls) which supports 14 extant native species. Just
95 seven extant species are native to both the upper and lower basins (Table 1; Wallace and
96 Zaroban 2013; Sigler and Zaroban 2018).

97 Aquatic species management in the Snake River basin has focused primarily on native trout.
98 Yellowstone Cutthroat Trout (including the fine-spotted form) are emphasized above Shoshone
99 Falls with Redband and Bull Trout emphasized below it (Figure 1). All are Species of Greatest
100 Conservation Need (SGCN) as well as listed Species of Special Concern in Idaho, Wyoming,
101 Nevada, or Oregon. Bull Trout are also federally listed as a threatened species under the United
102 States Endangered Species Act. Nevertheless, eight other SGCN species occur in the lower
103 Snake River basin (such as Leopard Dace *Rhinichthys falcatus*, Umatilla Dace *R. umatilla*, and
104 the Shoshone Sculpin *Cottus greenei*). Four non-salmonid SGCN species occur in the Upper
105 Snake River basin, including Bluehead Sucker *Catostomus discobolus* and Northern Leatherside
106 Chub *Lepidomeda copei* (Table 1), with only Mountain Whitefish *Prosopium williamsoni*
107 abundantly occurring in both lower and upper basins. In addition to fishes, the Upper Snake
108 River basin provides critical habitat for a plethora of other SGCN species. In addition to the
109 crayfish, fairy shrimp, pond snails, mud snails, mussels, frogs, and toads listed in Table 1, a wide
110 diversity of SGCN birds, mammals, reptiles, and invertebrates persist in the Upper Snake River
111 basin (IDFG 2017).

112 Identification of NFCAs that support at-risk fishes, as well as other aquatic species of
113 concern, has the potential to effectively integrate diverse conservation and management
114 actions within large watersheds across extensive landscapes. This includes the Upper Snake
115 River basin that has a diversity of ecosystems, land ownerships and uses, as well as at-risk fishes
116 and other aquatic species. For example, in the Upper Snake River basin, NFCAs provide an
117 opportunity to link the conservation of headwater Cutthroat Trout populations with
118 downstream habitats supporting non-game fishes such as Bluehead Sucker, Northern
119 Leatherside Chub, as well as conservation opportunities for other at-risk native aquatic species
120 ranging from amphibians to birds to mammals. Consequently, the potential benefits of NFCAs
121 are extremely diverse. For example, they may be managed to sustain the connectivity of critical
122 habitats at watershed scales. Connectivity serves to enhance population persistence by
123 facilitating natural metapopulation processes (Dunham and Rieman 1999; Hilderbrand and
124 Kershner 2000; Compton et al. 2008). Simultaneously, NFCAs may provide discrete hydrologic
125 units in which native fish and other aquatic communities can be isolated, if necessary, from
126 non-native invasions (Novinger and Rahel 2003; Fausch et al. 2009). To be most effective,
127 proposed NFCAs should be large enough to support natural landscape processes that
128 contribute to the long-term persistence of populations (Haak and Williams 2012). Yet NFCAs
129 may also be small enough to encourage the integration of substantive management actions by
130 diverse entities ranging from federal to tribal to state to private landholders. Consequently, the
131 establishment of NFCAs may facilitate more efficient and effective cooperative actions to
132 benefit a wider assemblage of native aquatic species.

133

134

Methods

135 We identified potential NFCAs through a process that ranked all subwatersheds
136 (Hydrological Unit Code [HUC] 12 watersheds) in the Upper Snake River basin based on native
137 trout abundance and distribution, modeled occurrence probabilities for native non-game fishes,
138 differential weighting of species based on their prevalence, drainage network connectivity, and
139 land protection status. Rankings were intended to identify watershed-scale areas from the
140 headwaters downstream where native trout overlap in distribution with, or occur near, native
141 non-game fishes and where watershed scale conservation would benefit entire fish
142 assemblages. Our analysis did not focus on identifying unique habitats with endemic fishes
143 (e.g., spring habitats with Shoshone Sculpin) or genetically unique subpopulations or subspecies
144 not currently recognized as a distinct species (e.g., Wood River Bridgelip Sucker or Big Lost
145 Mountain Whitefish), nor did we focus on large river fishes (e.g., White Sturgeon *Acipenser*
146 *transmontanus*) or the mainstem Snake River because of the difficulty in managing large rivers
147 to their headwaters per the NFCA concept. Clusters of high ranking subwatersheds (i.e., the top
148 25%) were aggregated and characterized based on the native fish assemblage, land ownership
149 and protected status, watershed size, habitat conditions, and future threats.

150

Fish Data

151 Many different data sources were used to define the distribution and abundance of native
152 trout and the occurrence of native non-game fishes. Distribution and abundance data for native
153 trout were primarily derived from range-wide assessment databases. Yellowstone Cutthroat
154

155 Trout data (including the fine-spotted form) were based on the 2010 range-wide assessment
156 database (Gresswell 2011), and Redband Trout data were based on the 2012 range-wide
157 assessment database (Muhlfeld et al. 2015). For each database, only conservation populations
158 were used to define distributions (Figure 1), and abundance was based on the midpoint of
159 categorical population abundances in the database (e.g., 0 – 35, 35 – 100, 101 – 250, 251 – 625,
160 625 – 1250, >1250 fish per km). Conservation populations were defined as those populations
161 that had <10% genetic introgression or had unique genetic, ecological or behavioral attributes
162 (e.g., adfluvial behavior) (UDWR 2000; Gresswell 2011; Muhlfeld et al. 2015). Bull Trout
163 distribution data were obtained from Streamnet, and abundance data from agency databases
164 (see Acknowledgements).

165 Native non-game fishes data were assembled from fish collections made across the Upper
166 Snake River basin, and those data were used to develop species probability of occurrence
167 models. Non-game fishes were sampled primarily by electrofishing but other methods were
168 also used; for example, non-game species in Idaho were primarily sampled during Idaho
169 Department of Fish and Game (IDFG) and Idaho Department of Environmental Quality
170 electrofishing surveys (Meyer et al. 2013) while both electrofishing and minnow traps were
171 used in other datasets (Blakney 2012). In all, over 3047 fish collection records were used to
172 determine species occurrences (see Acknowledgments for sources). From these collections,
173 presence-absence data were used to develop species-specific species distribution models
174 (random forest; Breiman 2001) where probabilities of occurrence were modeled as a function
175 of multiple environmental variables (% converted land, canal density, etc.); see supplementary
176 files for details. The models for all species, except Leopard Dace, indicated good predictive
177 ability with 10-fold cross-validated AUC values >0.75. The model for Leopard Dace, a species
178 with only 10 occurrences, had the poorest predictive ability (AUC = 0.695). Species-specific
179 models were then used to predict probability of occurrence in perennial stream segments in
180 the National Hydrography Dataset Plus (Version 1; 1:100,000 scale) with drainage areas larger
181 than 159,100 km². Predictions were only made in sub-basins within the probable native ranges
182 sampled by Meyer et al. (2013) and Gamett (2003).

183

184 *Watershed Rankings*

185 To identify potential NFCAs, every subwatershed in the Upper Snake River basin was ranked
186 based on native trout distribution and abundance data, non-game fish probabilities of
187 occurrence data, river network connectivity, and the percent of subwatersheds encumbered in
188 Gap Analysis Program (GAP) Status 1 or 2 protected lands (e.g., wilderness, or national parks)
189 (USGS 2011). The analysis was constructed to generally give higher ranks to clusters of inter-
190 connected subwatersheds with abundant trout populations and high native fish richness (high
191 biodiversity) within or near protected lands. More specifically, subwatershed rankings were
192 obtained using the Additive Benefit Function in Zonation v3.0 conservation planning software
193 to account for river system connectivity (Moilanen 2007; Moilanen et al. 2008; Moilanen et al.
194 2011). Zonation produces a hierarchical ranking of all subwatersheds (rescaled from 0 to 100)
195 based on the minimum marginal loss across species-specific input values given species weights
196 (minimum biodiversity loss) and offset by a cost function: $\delta_i = 1/c_i w_j \sum_j \Delta V_j$ where δ_i = the
197 marginal loss across all j species for subwatershed i , $c_i = 100 - \% \text{ subwatershed protected}$,
198 where % protected was based on protected lands identified as GAP status 1 or 2 lands (w_j = the

199 weight for species j based on professional judgment for native trouts or non-game species
200 prevalence – one measure of rarity - defined by Meyer et al. (2013)(see Table 1); and ΔV_j is
201 the marginal loss of species j values between all remaining subwatersheds minus the value
202 within subwatershed i (see Moilanen et al. 2011). The value of V differed by species. For
203 conservation populations of Yellowstone Cutthroat Trout and Redband Trout (see above) V =
204 the midpoint of fish per km ranges reported for those populations, V = subwatershed fish per
205 km average (from fish survey data) for the current distribution of Bull Trout (a value of 1 was
206 used if no abundance data were available), and V = the probability of occurrence (range from 0
207 to 1) for all non-game species. Rankings were determined by computing the minimum marginal
208 loss across all species for each subwatershed, and removing the subwatershed with the lowest
209 marginal loss. This process effectively removed the subwatershed from the landscape that
210 resulted in the minimum biodiversity loss given species weights and amount of protected lands
211 in the entire Upper Snake River basin. The removal process was repeated iteratively until only
212 one subwatershed remained during the last iteration, that is, the subwatershed with the
213 highest marginal loss across all fish species and the most important subwatershed from a
214 biodiversity standpoint. The sequence of removal resulted in the subwatershed rank. Trout data
215 were represented by the spatial hydrography framework for each data source, and the
216 probability of occurrence for non-game species was attributed on NHDPlus. All data were then
217 converted to a 300-m grid for the analysis.

218 River networks are a nested hierarchy of drainage systems that pose challenges to
219 conservation of fishes residing in riverine environments. Consequently, NFCAs are watersheds
220 managed in their entirety for native aquatic communities. For this reason, river network
221 connectivity was used to impart a penalty on the marginal loss across species based on whether
222 a neighboring subwatershed has already been removed during the ranking process (i.e., has a
223 lower rank). The penalty specifically translated into a reduction in subwatershed value (δ_i)
224 based on the proportion of subwatersheds that had been removed upstream or downstream of
225 the focal subwatershed (see Section 2.4.4 in Moilanen et al. 2011). The same penalties were
226 used for all 21 species and were strong if upstream subwatersheds had already been removed
227 and weak if downstream subwatersheds had been removed. Although connectivity was
228 interrupted by large dams (i.e., those with reservoirs with ≥ 4 km² surface area), smaller dams or
229 other barriers were not used to break connectivity because they are much more capable of
230 being managed for fish passage. Figure 2 illustrates the analytical process.

231 232 *Potential NFCAs*

233 Clusters of subwatersheds representing independent drainage networks within the top 25%
234 of the landscape (rank >75) were then aggregated into potential NFCAs. We summarized
235 selected attributes of potential NFCAs: mean subwatershed rank; watershed size; documented
236 native fish occurrences; presence of non-game fishes of greatest conservation need; percent of
237 watershed protected; percent of perennial stream corridor protected; habitat integrity of
238 subwatershed; and future security of subwatershed. The mean rank of subwatersheds in an
239 NFCAs was computed as an area-weighted mean from the subwatershed ranking analysis.
240 Documented species occurrences were determined from range-wide assessment databases for
241 native trout and documented presence of native non-game species in fisheries surveys. State
242 wildlife action plans were used to identify Species of Greatest Conservation Need in each state

243 (e.g., IDFG 2017). Land protection status was based on GAP Status 1 and 2 lands (USGS 2011).
244 Subwatershed habitat integrity was based on the Trout Unlimited Conservation Success Index; a
245 composite index ranging from 5 (low integrity) to 25 (high integrity) based on indicators of land
246 protection, watershed connectivity, watershed condition, water quality, and flow regime scored
247 each from 1 (low integrity) to 5 (high integrity) (Williams et al. 2007). Future security of
248 subwatersheds was also based on the Trout Unlimited Conservation Success Index where future
249 security is an index ranging from 5 (low security) to 25 (high security) based on individuals
250 scores (1 to 5) for land conversion, resource extraction, energy development, climate change,
251 and introduced species (Williams et al. 2007).

252

253

Results

254 Watershed rankings from the Zonation analysis identified entire subbasins where all
255 subwatersheds ranked high as well as individual or small clusters of a few high-ranking
256 subwatersheds (Figure 3). Potential NFCA watersheds were primarily clustered in headwater
257 areas, both within individual river systems (e.g., Boise, Payette, Little Jacks, Little Lost, Goose
258 drainages), and across the Snake River as a whole (e.g., South Fork Snake, Salt, Blackfoot). The
259 Payette, Boise, South Fork Snake, Portneuf, and Blackfoot river subbasins represented large
260 clusters of subwatersheds with ranks exceeding 75 (of a range from 0 to 100), and representing
261 the top 25% of the entire Upper Snake River basin. Smaller aggregations were present in the
262 Upper Malheur River, Upper Jarbidge River, Goose Creek, Little Lost River, and Fall
263 River/Conant Creek (Henrys Fork watershed). Examples of individual drainages with one or a
264 few high ranking subwatersheds were: Little Jacks Creek, Jack Creek (NV), Indian Creek, Cassia
265 Creek, Upper Raft River, and Bitch Creek (Figure 4A).

266 After the clusters of high ranking subwatersheds (top 25%) were aggregated, 44 watersheds
267 (range from 56 to 4,344 km²) were identified as potential NFCAs based on their high rank and
268 native species assemblages (Table 2). All potential NFCAs supported at least one native trout
269 species, and all but three supported at least one native non-game species. Of those, 13
270 watersheds supported at least one non-game species of greatest state conservation need
271 (Table 2; Figure 4A).

272 Land status (USGS 2004) within watersheds ranged from 72% private land (Portneuf River)
273 to nearly the entire upper watershed protected in public land and national parks (Upper Snake
274 River; Cottonwood Creek; Fall River) (Table 2; Figure 4A). Although stream corridors ranged
275 widely in level of protection (Figure 4B), land status was not a reliable predictor of habitat
276 integrity or future security.

277 Habitat integrity scores ranged from 12 to 25 (Table 2). Habitat integrity in the Upper Snake
278 River basin was low in areas with extensive agricultural or urban land use, and high in upper
279 elevation mountainous regions (Figure 5A). Watersheds comprised largely of public land (>90%)
280 exhibited habitat integrity scores ranging from the highest possible (25) to 12. In potential
281 NFCAs, habitat integrity was high for Pacific Creek in Grand Teton National Park (score 25 out of
282 25). In contrast, despite exhibiting species-rich fish assemblages (Table 2), habitat integrity was
283 low for the Portneuf River and Conant Creeks (scores of 12 out of 25) due to low scores (1 out
284 of 5) in each of five individual indicators in at least one subwatershed. While increased levels of
285 private land ownership within a watershed generally resulted in lower habitat integrity scores,

286 of the eleven watersheds with >35% private land ownership, habitat integrity scores ranged
287 from 12 in the Portneuf River and Conant Creek (Henry's Fork subbasin) to 18 in the Upper Raft
288 River, where 70% of the watershed is in private ownership.

289 Likewise, future security of habitats (and fishes) was variable across the basin (Figure 5B). It
290 was typically high in protected areas and in areas with low human development. Within
291 potential NFCAs, future security was high within several watersheds in national parks (scores 21
292 out of 25), whereas Indian Creek in Hells Canyon had low future security (scores 11 out of 25)
293 because of threats from land conversion, resource extraction, and climate change (Table 2).
294 Bear Creek in the South Fork Snake River subbasin had a future security score of 13, despite
295 being 99% public lands and a habitat integrity score of 22, while Canyon Creek in the Henry's
296 Fork Snake River subbasin, the Lower South Fork Snake River, and several Payette River
297 tributaries all had future security scores of 12 and 13 (Table 2).

298 Identification of Potential NFCAs highlighted native trout distributions and abundance
299 coupled with presence of native non-game species, particularly rare and sensitive species (i.e.,
300 Species of Greatest Conservation Need) (Figure 4). Potential NFCAs in the lower Snake River
301 basin included assemblages of Redband and Bull trout and from one to eleven non-game
302 species (except for Indian Creek which contained native trout only). Non-game SGCN species (n
303 = 4) are disproportionately represented in potential NFCAs in the upper basin.

304 To illustrate differences in the characteristics of potential NFCAs, we contrasted the
305 characteristics of three watersheds: Jarbidge River, Goose Creek, and Upper Blackfoot River
306 (Figure 6). The Jarbidge River supports two native trouts, Redband Trout and Bull Trout,
307 whereas the others have only Yellowstone Cutthroat Trout. The watersheds ranged in size from
308 878 to 1,842 km², but only two, Goose Creek and Upper Blackfoot River, have natural
309 downstream extents defined by dams impounding reservoirs. The Jarbidge River has no
310 definitive downstream boundary and is connected to downstream rivers (a dam isolates the
311 Bruneau River from the Snake River much further downstream). The amount of public land in
312 the watersheds ranges from 56 to 90%, with USDA Forest Service lands in the headwaters and a
313 mix of Bureau of Land Management and state lands at lower elevations. Perennial stream
314 corridors were 72 to 87% public land except in the Upper Blackfoot River where 41% of stream
315 kilometers were on public land; however, the mainstems of all streams and rivers were largely
316 on private land. Little of the perennial streams had formal land protections, except in the
317 Jarbidge River where 37% of perennial stream kilometers have formal protections in wilderness
318 or wild and scenic river designation. Habitat integrity and future security for these three
319 watersheds were moderate to high, in contrast for example to the future security of the South
320 Fork Snake River and tributaries, which are lower and threatened by introduced species (e.g.,
321 Rainbow Trout), climate change (high drought risk), and energy development (potential
322 hydropower development).

323

324

Discussion

325 Aquatic habitat in the western United States has been degraded by myriad factors
326 including: over-grazing by domestic livestock (Beschta et al. 2013), water withdrawal (Deacon et
327 al. 2007), oil and gas development (Dauwalter 2013), urbanization (Williams et al. 2007),
328 hydroelectric development (Thurow et al. 2000), and forestry and mining practices (Lee et al.

329 1997); among other anthropogenic impacts. In turn, the distributions of inland trout have been
330 substantially reduced; for example, all inland Cutthroat Trout subspecies occupy less than half
331 of their historical habitat (Thurow et al. 1997; Haak and Williams 2013). Salmon have been
332 extirpated from some rivers (Praggastis and Williams 2013), and non-game fishes have
333 experienced declines in abundance and distribution in western North America (Brouder and
334 Scheurer 2007). Habitat conditions are expected to continue to change in the future as runoff
335 timing, stream discharge, and water temperatures are altered by climate warming (Wenger et
336 al. 2011; Meyer et al. 2014a; Beschta et al. 2013; Isaak et al. 2015).

337

338 *Benefits of NFCAs for Native Fishes*

339 Native Fish Conservation Areas provide both a conceptual and formal framework for
340 collaborative, watershed-scale conservation and restoration efforts targeting broader native
341 fish communities that may include native trout in headwaters and native non-game species in
342 both headwaters and downstream areas in western river systems (Williams et al. 2011).
343 Watershed scale conservation and restoration can improve instream habitat diversity closely
344 linked to fish diversity in the Upper Snake River basin (Walrath et al. 2016). Beaver are
345 increasingly important for restoring stream channels incised by livestock grazing and other
346 impacts. Such efforts can be critical for SCGN species like Northern Leatherside Chub that have
347 higher prevalence at sites with complex streamflows associated with beaver dams (Dauwalter
348 and Walrath 2018). Stream habitat restoration and increased connectivity at the watershed
349 scale is expected to improve current conditions for native aquatic assemblages and increase
350 their resiliency to future threats like climate change (Thurow et al. 1997; Zoellick et al. 2005;
351 Isaak et al. 2015).

352 We used a quantitative, data-driven approach to identify key locations in the Upper Snake
353 River basin where watershed-scale restoration efforts will benefit native fish communities, such
354 as in Goose Creek, the Upper Blackfoot River, and Conant Creek (Figure 4A). Watershed-scale
355 restoration might also be a management priority in watersheds that are juxtaposed to lands
356 currently designated for protection as Wild and Scenic Rivers, Wilderness, or National Parks
357 because of the additional future security for the watersheds and aquatic communities.
358 Examples of potential NFCA watersheds in designated status include the Upper South Fork
359 Snake, Pacific Creek, and Hoback Creek (Table 2; Figure 4A). Ultimately, adoption of these
360 potential NFCAs as functional NFCAs will require a stakeholder-driven, collaborative process
361 ensuring feasibility assessment and implementation (Dauwalter et al. 2011; Birdsong et al.
362 2015).

363 Decades ago, Lee et al. (1997) anticipated the need for NFCAs after examining historical and
364 contemporary distributions of 15 native salmonid taxa in the interior Columbia River basin and
365 portions of the Klamath River and Great basins. The authors evaluated the native salmonids we
366 examined in this study (Redband Trout, two forms of Yellowstone Cutthroat Trout, Bull Trout
367 and Mountain Whitefish) (Lee et al. 1997; Thurow et al. 1997). Their work identified habitat and
368 biodiversity strongholds for native salmonids in the northern Cascades, the central Idaho
369 mountains, and the Snake River headwaters and their connecting river corridors. Thurow et al.
370 (1997) noted that maintaining the integrity of these declining native fishes and aquatic systems
371 for the long term would require a network of well-connected, high-quality habitats
372 (strongholds) that supported a diverse assemblage of native species including a full expression

373 of their life histories. Our finer-scaled analysis of the Upper Snake River basin further
374 illuminates areas of diverse, high-quality, interconnected habitat in the headwaters of the
375 Boise, Payette, and Upper Snake river drainages and the potential for NFCAs in several of these
376 same watersheds (Figures 4 and 6).

377 For the NFCA approach to function and assist both the public interest and management
378 agencies' long-term planning efforts, it is essential that NFCAs be identified at a scale that is
379 practical and manageable. Aggregating smaller subwatersheds with common species
380 assemblages and management concerns into larger management units (as potential NFCAs)
381 ensures NFCA designation will be an effective management tool (Birdsong et al. 2019, Chapter
382 X). Although our unit of analysis was a subwatershed (~12,000 hectares), our analysis also
383 explicitly accounted for connectivity of adjacent subwatersheds. This facilitated aggregation
384 into larger watersheds where management considerations and actions were likely to be
385 consistent. For example, IDFG manages Yellowstone Cutthroat Trout at the subbasin level (IDFG
386 2007), a scale representing a natural way to aggregate our results into larger units, as we
387 illustrate (Figure 4).

388 Watershed scale NFCAs are also compatible with the proposed management strategies of
389 many state and federal natural resource management plans, because they rely on identifying
390 management units based on habitat conditions, genetics, and population status. IDFG's
391 Management Plan for conservation of Yellowstone Cutthroat Trout (IDFG 2007) defines 13
392 Geographic Management Units addressing abundance, trends, genetics, and an evaluation of
393 existing threats (Meyer et al. 2006). The status of Redband Trout populations is summarized in
394 a similar manner below Shoshone Falls (Meyer et al. 2014b).

395 The NFCA concept can also be used as an organizational framework for cross-partnership
396 collaborations focusing on strategic restoration of fish habitat. Several multi-agency
397 partnerships, such as the Western Native Trout Initiative (WNTI;
398 <http://www.westernnativetrout.org>) and the Desert Fish Habitat Partnership (DFHP;
399 www.desertfhp.org), are collaborative, multi-agency and multi-state partnerships focused on
400 native trout conservation and restoration across the western United States and conservation
401 and restoration of habitats used by non-salmonid desert fishes, respectively. The WNTI and
402 DFHP partnerships commonly collaborate on habitat restoration projects (Dauwalter et al.
403 2019, Chapter X). The NFCA concept provides an umbrella framework guiding expanded
404 collaboration of watershed-scale restoration efforts that could benefit native trout in
405 headwater streams as well as native, non-game fishes in downstream, warmer mainstem
406 habitats. For example, our Upper Snake River analysis informs identification of focal watersheds
407 for such cross-partnership collaboration, planning, and fund-raising (e.g., Dauwalter et al. 2011;
408 Haak and Williams 2013; Dauwalter et al. 2019, Chapter X).

409 Native sport fishes such as Cutthroat Trout and Black Basses *Micropterus spp.* have been
410 proposed as key species defining NFCAs because they range widely, are relatively well known
411 compared to many other native stream fishes, and are of primary interest to both recreational
412 anglers and state and federal management agencies; yet they are relatively sensitive to
413 disturbance (Williams et al. 2011; Birdsong et al. 2019, Chapter X). Restoring and reconnecting
414 stream networks has been proposed as a method to conserve native trout in the face of climate
415 change by creating larger stronghold populations and facilitating development of migratory life
416 histories that aid in long-term population persistence (Haak and Williams 2013). Haak and

417 Williams (2013) describe such efforts for Rio Grande Cutthroat Trout *O. c. virginialis*, including
418 restoration of 240 km of interconnected habitats in the Rio Costilla area of New Mexico that
419 also creates a refuge for other native fishes of the Rio Grande Basin.

420 The NFCA concept has also been applied to explore watershed level conservation and
421 stakeholder partnerships in Colorado and across the Southeast U.S. Dauwalter et al. (2011)
422 compared the distribution of remaining Colorado River Cutthroat Trout *O. c. pleuriticus* with
423 known distributions of three sensitive warmwater fishes (Roundtail Chub *Gila robusta*,
424 Bluehead Sucker, and Flannelmouth Sucker *Catostomus latipinnis*) to identify potential NFCAs in
425 the Upper Colorado River drainages of Colorado, Utah, and Wyoming. Several watershed-scale
426 opportunities were identified, including Muddy Creek in the Little Snake River, where multiple
427 agencies and non-profit groups are working to reconnect headwaters with downstream reaches
428 in order to conserve the entire native fish community (Compton et al. 2008).

429 In the Southeast U.S., the NFCA concept has been proposed as a preferred method to
430 protect native stream fishes. Birdsong et al. (2015) recommended the use of the NFCA concept
431 to protect native Black Basses, which are keystone species in southeastern stream systems.
432 Their protection in large, functional watersheds would simultaneously help conserve numerous
433 less well known, native stream fishes in the region. Efforts are underway to identify NFCA
434 watersheds for keystone Black Bass species including: Guadalupe Bass *Micropterus treculii* in
435 the Llano River watershed in central Texas; Redeye Bass *M. coosae* in South Carolina's upper
436 Savannah River watershed; and Shoal Bass *M. cataractae* in the Chipola River watershed in
437 north-central Florida.

438

439 *Benefits of NFCAs for Other Aquatic Species*

440 Management focused on resilient watersheds will also likely benefit other imperiled aquatic
441 species such spotted frogs *Rana luteiventris*, western ridged mussel *Gonidea angulata*, other
442 diverse SCGN species (Table 1), as well as riparian obligate species such as the sage grouse
443 *Centrocercus urophasianus* that require wet meadow habitat for early juvenile rearing. Such an
444 approach would engage multiple partners and potentially leverage increased funding sources in
445 a more efficient manner to serve a wide diversity of species (Birdsong et al. 2019, Chapter X).

446 An example of this multi-organization partnership approach was implemented in 2015 on
447 North Carolina's Little Tennessee River (LTR), which became the nation's first designated NFCA
448 (Harris and Williams 2016; Leslie et al. 2019, Chapter X, this volume). The LTR is an important
449 global biological hotspot and is host to a unique assemblage of fish, amphibians, mollusks,
450 crayfish, and aquatic insects. The North Carolina Wildlife Federation gathered interested
451 organizations, agencies, and business to discuss long-term conservation of the LTR. As a result,
452 the LTR Native Fish Conservation Partnership formed to assist management of the LTR
453 watershed as an NFCA. The LTR Native Fish Conservation Partnership includes 25 collaborative
454 partners comprising non-governmental organizations, federal and state agencies, the Eastern
455 Band of Cherokee Indians, and several private businesses.

456

457 *Collaborative Benefit of NFCAs*

458 Addressing specific local and regional aquatic conservation issues through the NFCA concept
459 can facilitate important socio-ecological perspectives that are key to achieving both biological
460 goals and the social and management goals of NFCA partnership members (e.g., the Little

461 Tennessee River NFCA). Establishing the multi-stakeholder partnership, required to support an
462 NFCA, fosters a collaborative, interdisciplinary approach necessitating active dialogue for
463 sharing values and perspectives. NFCA collaboration provides the opportunity for effectively
464 discussing and resolving conflict among user groups, managers, and fisheries biologists (Hand et
465 al. 2018; Birdsong et al. 2019 Chapter X, this volume). Rieman et al (2000) used a broad-scale
466 and spatially explicit classification of subbasins in the Northwest to examine opportunities and
467 conflicts related to terrestrial and aquatic ecosystem management. Birdsong and colleagues
468 (2019, Chapter X, this volume) discuss a 6-year multi-partner effort across watersheds in Texas
469 that led to the development of the Texas Native Fish Conservation Areas Network. The Texas
470 NFCA Network protects more than 90 freshwater fishes considered species of greatest
471 conservation need and their associated watersheds through designation of twenty Native Fish
472 Conservation Areas identified to preserve the unique Texas freshwater fish diversity.

473 Because human disturbance is strongly associated with the condition of both aquatic and
474 terrestrial ecosystems, the commonality in management goals and opportunities challenges
475 managers of multiple disciplines to work together. Spatially explicit classifications such as
476 NFCAs provides a mechanism for better integrating management. Rieman et al. (2000)
477 suggested that such approaches assist integration through: (1) communication among
478 disciplines; (2) effective prioritization of limited conservation and restoration resources; and (3)
479 establishing a framework for experimentation and demonstration of restoration techniques.
480 Establishing a multi-disciplinary and multi-organization partnership to support NFCAs is a critical
481 element for creating the public and institutional support necessary for more effective, long-
482 term conservation and restoration of aquatic systems, as demonstrated eloquently in the Little
483 Tennessee River (Leslie et al. 2019) and across the Texas NFCA Network (Birdsong et al. 2019).

484

485

Acknowledgments

486 We thank the following individuals for sharing Snake River basin fish population data: K.
487 Meyer, IDFG; B. Gamett, Salmon-Challis National Forest; D. Miller, Wyoming Game and Fish
488 Department; B. Bangs, Oregon Department of Fish and Wildlife; J. Pappani, Idaho Department
489 of Environmental Quality; R. Hughes, U.S. Environmental Protection Agency; E. Keeley, Idaho
490 State University; J. DeRito, Henrys Fork Foundation(HFF)/Trout Unlimited; M. Lien, Friends of
491 the Teton River; and contributors to the HFF database: BLM, USFS, WGFD, IDFG, and FTR. We
492 thank those who responded to our survey regarding the characterization of NFCAs. We also
493 thank S. Grunder and J. Dillon (IDFG), S. Hoefler and A. Berglund (BLM), R. Perkins (ODFW), M.
494 Lien and A. Vebeten (Friends of the Teton River), L. Mabey (Caribou-Targhee National Forest),
495 D. Miller (WGFD) for assisting us in reviewing the data and analyses. Funding for this work was
496 provided by the National Fish and Wildlife Foundation and Patagonia’s Environmental Fund and
497 World Trout Initiative.

498

499

Supplementary Files

500 www.tu.org/USRB-multisp-assmt

501

502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543

References

Benke, A. C. and C. E. Cushing, C.E., editors. 2005. Rivers of North America. Academic Press, Burlington, Massachusetts.

Beschta, R. L., D. L. Donahue, D. A. DellaSala, J. J. Rhodes, J. R. Karr, M. H. O'Brien, T. L. Fleischner, C. Deacon Williams. 2013. Adapting to climate change on western public lands: addressing the ecological effects of domestic, wild, and feral ungulates. *Environmental Management* 51:474-491.

Birdsong, T. W., M. S. Allen, J. E. Claussen, G. P. Garrett, T. B. Grabowski, J. Graham, F. Harris, A. Hartzog, D. Hendrickson, R. A. Krause, J. Leitner, J. M. Long, C. K. Metcalf, D. P. Phillipp, W. F. Porak, S. Robinson, S. M. Sammons, S. Shaw, J. E. Slaughter, and M. D. Tringali. 2015. Native Black Bass Initiative: implementing watershed-scale approaches to conservation of endemic black bass and other native fishes in the southern United States. *American Fisheries Society Symposium* 82:363-378.

Birdsong, T. W., G. P. Garrett, B. J. Labay, M. Bean, P. Bean, J. Botros, M. Casarez, A. Cohen, T. Heger, A. Kalmbach, D. Hendrickson, S. Magnelia, K. Mayes, M. McGarrity, R. McGillicuddy, M. Parker, and S. Robertson. 2019. Texas Native Fish Conservation Areas Network: strategic investments in restoration and preservation of freshwater fish diversity. Pages XXX - XXX in D. C. Dauwalter, T. W. Birdsong, and G. P. Garrett, editors. *Multispecies and watershed approaches to freshwater fish conservation*. American Fisheries Society, Bethesda, Maryland.

Blakney, J. R. 2012. Historical connectivity and contemporary isolation: population genetic structure of a rare high-desert minnow, the northern leatherside chub (*Lepidomeda copei*). M.S. Idaho State University, Pocatello.

Breiman, L. 2001. Random forests. *Machine Learning* 45:5-32.

Brouder, M. J., and J. A. Scheurer, editors. 2007. Status, distribution, and conservation of native freshwater fishes of western North America. *American Fisheries Society Symposium* 53.

Campbell, M. R., C. C. Kozfkay, K. A. Meyer, M. S. Powell, and R. N. Williams. 2011. Historical influences of volcanism and glaciation in shaping mitochondrial DNA variation and distribution of Yellowstone cutthroat trout across its native range. *Transactions of the American Fisheries Society* 140:91-107.

Compton, R. I., W. A. Hubert, F. J. Rahel, M. C. Quist, and M. R. Bower. 2008. Influences of fragmentation on three species of native warmwater fishes in a Colorado River Basin headwater stream system. *North American Journal of Fisheries Management* 28:1733-1743.

544 Dauwalter, D. C., J. S. Sanderson, J. E. Williams, and J. R. Sedell. 2011. Identification and
545 implementation of Native Fish Conservation Areas in the Upper Colorado River Basin.
546 Fisheries 36:278-288.
547

548 Dauwalter, D. C. 2013. Fish assemblage associations and thresholds with existing and projected
549 oil and gas development. Fisheries Management and Ecology 20:289-301.
550

551 Dauwalter, D. C., and J. D. Walrath. 2018. Beaver dams, streamflow complexity, and the
552 distribution of a rare minnow, *Lepidomeda copei*. Ecology of Freshwater Fish 2018:27:606–
553 616.
554

555 Dauwalter, D. C., S. Vail-Muse, T. Thompson, K. Johnson, and J. Whittier. 2019. Partnering on
556 multispecies aquatic assessments to inform efficient conservation delivery. Pages XXX - XXX
557 in D. C. Dauwalter, T. W. Birdsong, and G. P. Garrett, editors. Multispecies and watershed
558 approaches to freshwater fish conservation. American Fisheries Society, Bethesda,
559 Maryland.
560

561 Deacon, J. E., A. E. Williams, C. D. Williams, and J. E. Williams. 2007. Fueling population growth
562 in Las Vegas: how large-scale groundwater withdrawal could burn regional biodiversity.
563 BioScience 57:688-698.
564

565 Dunham, J. B., and B. E. Rieman. 1999. Metapopulation structure of bull trout: influences of
566 physical, biotic and geometrical landscape characteristics. Ecological Applications 9:642-
567 655.
568

569 Fausch, K. D., B. E. Rieman, J. B. Dunham, M. K. Young, and D. P. Peterson. 2009. Invasion versus
570 isolation: trade-offs in managing native salmonids with barriers to upstream movement.
571 Conservation Biology 23:859-870.
572

573 Gamett, B. L. 2003. A summary: the origin of fishes in the Sinks Drainages of southeastern
574 Idaho. Page 16 in R. W. Van Kirk, J. M. Capurso, and B. L. Gamett, editors. The Sinks
575 symposium: exploring the origin and management of fishes in the Sinks Drainages of
576 southeastern Idaho. Idaho Chapter American Fisheries Society, Boise.
577

578 Gresswell, R. E. 2011. Biology, status, and management of the Yellowstone cutthroat trout.
579 North American Journal of Fisheries Management 31:782-812.
580

581 Haak, A. L., and J. E. Williams. 2012. Spreading the risk: native trout management in a warmer
582 and less-certain future. North American Journal of Fisheries Management 32:387-401.
583

584 Haak, A. L., and J. E. Williams. 2013. Using native trout restoration to jumpstart freshwater
585 conservation planning in the Interior West. Journal of Conservation Planning 9:38-52.
586

587 Hand, B. K., C.G. Flint, C. A Frissell, C. C. Muhlfeld, S. P. Devlin, B. P. Kennedy, R. L. Crabtree, W.
588 A. McKee, G. Luikart, and J. A. Stanford. 2018. A social-ecological perspective for riverscape
589 management in the Columbia River Basin. *Frontiers in Ecology and Environment* 16(S1):
590 S23-S33.
591

592 Harris, F. and R. N. Williams. 2016. Introducing No. 1: Little Tennessee River Basin Becomes First
593 Native Fish Conservation Area. *Flyfisher*. Spring Summer 2016.
594

595 Hilderbrand, R. H., and J. L. Kershner. 2000. Conserving inland cutthroat trout in small streams:
596 how much stream is enough? *North American Journal of Fisheries Management* 20:513-520.
597

598 IDFG (Idaho Department of Fish and Game). 2007. Management plan for conservation of
599 Yellowstone cutthroat trout in Idaho. Idaho Department of Fish and Game, Technical
600 Report, Boise, Idaho. 118 pp.
601

602 IDFG (Idaho Department of Fish and Game). 2017. Idaho State Wildlife Action Plan, 2015. Idaho
603 Department of Fish and Game. Grant No.: F14AF01068 Amendment, Sponsored by the US
604 Fish and Wildlife Service, Wildlife and Sport Fish Restoration Program. #1. Boise, Idaho
605 Available from: <http://fishandgame.idaho.gov/>.
606

607 Isaak, D. J., M. K. Young, D. E. Nagel, D. L. Horan, and M. C. Groce. 2015. The cold-water climate
608 shield: delineating refugia for preserving salmonid fishes through the 21st century. *Global
609 Change Biology* 21:2540-2553.
610

611 Jelks, H. L., S. J. Walsh, S. Contreras-Balderas, E. Díaz-Pardo, N. M. Burkhead, D. A. Hendrickson,
612 J. Lyons, N. E. Mandrak, F. McCormick, J. S. Nelson, S. P. Platania, B. A. Porter, C. B. Renaud,
613 J. J. Schmitter-Soto, E. B. Taylor, and M. L. Warren. 2008. Conservation status of imperiled
614 North American freshwater and diadromous fishes. *Fisheries* 33:372-407.
615

616 Lee, D. C., J. R. Sedell, B. E. Rieman, R. F. Thurow, and J. E. Williams. 1997. Broadscale
617 assessment of aquatic species and habitats. Pages 1058–1496 in T. M. Quigley and S. J.
618 Arbelbide, editors. An assessment of ecosystem components in the Interior Columbia Basin
619 and portions of the Klamath and Great Basins. U.S. Department of Agriculture Forest
620 Service, PNW-GTR-405, Portland, Oregon.
621

622 Leslie, A., E. McCombs, and F. Harris. 2019. Little Tennessee River Basin Native Fish
623 Conservation Partnership: aquatic conservation on a landscape scale. Pages XXX - XXX in D.
624 C. Dauwalter, T. W. Birdsong, and G. P. Garrett, editors. Multispecies and watershed
625 approaches to freshwater fish conservation. American Fisheries Society, Bethesda,
626 Maryland.
627

628 Master, L. L., B. A. Stein, L. S. Kutner, and G. A. Hammerson. 2000. Vanishing assets:
629 conservation status of U.S. species. Pages 93– 118 in B. A. Stein, L. S. Kutner, and S. J.

630 Adams, editors. Precious heritage: the status of biodiversity in the United States. Oxford
631 University Press, New York.

632

633 Meyer, K. A., D. J. Schill, and J. A. Lamansky, Jr. 2006. Status of Yellowstone cutthroat trout in
634 Idaho. *Transactions of the American Fisheries Society* 135:1329-1347.

635

636 Meyer, K. A., J. A. Lamansky, Jr., D. J. Schill, and D. W. Zaroban. 2013. Nongame fish species
637 distribution and habitat associations in the Snake River Basin of southern Idaho. *Western*
638 *North American Naturalist* 73:20-34.

639

640 Meyer, K. A., E. I. Larson, C. L. Sullivan, and B. High. 2014a. Population trends of Yellowstone
641 cutthroat trout and nonnative trout in Idaho's upper Snake River basin. *Journal of Fish and*
642 *Wildlife Management* 5:227-242.

643

644 Meyer, K. A., D. J. Schill, E. R. J. M. Mamer, C. C. Kozfkay, and M. R. Campbell. 2014b. Status of
645 redband trout in the Upper Snake River Basin of Idaho. *North American Journal of Fisheries*
646 *Management* 34:507-523.

647

648 Moilanen, A. 2007. Landscape zonation, benefit functions and target-based planning: unifying
649 reserve selection strategies. *Biological Conservation* 134:571-579.

650

651 Moilanen, A., J. Leathwick, and J. Elith. 2008. A method for spatial freshwater conservation
652 prioritization. *Freshwater Biology* 53:577-592.

653

654 Moilanen, A., L. Meller, J. Leppänen, A. Arponen, and H. Kujala. 2011. Zonation: spatial
655 conservation planning framework and software. version 3.0, University of Helsinki, Finland.

656

657 Muhlfeld, C. C., S. E. Albeke, S. L. Gunckel, B. J. Writer, B. B. Shepard, and B. E. May. 2015.
658 Status and conservation of Interior Redband Trout in the western United States. *North*
659 *American Journal of Fisheries Management* 35:31-53.

660

661 Novinger, D. C., and F. J. Rahel. 2003. Isolation management with artificial barriers as a
662 conservation strategy for cutthroat trout in headwater streams. *Conservation Biology*
663 17:772-781.

664

665 Praggastis, A., and J. E. Williams. 2013. Salmon's presence in Nevada's past. *Nevada Historical*
666 *Society Quarterly* 56(1-2):14-32.

667

668 Rieman, B. E., D. C. Lee, R. F. Thurow, P. F. Hessburg, and J. R. Sedell. 2000. Toward an
669 integrated classification of ecosystems: defining opportunities for managing fish and
670 forest health. *Environmental Management* 25:425-444.

671

672 Roux, D. J., J. L. Nel, P. J. Ashton, A. R. Deacon, F. C. de Moor, D. Hardwick, L. Hill, C. J.
673 Kleynhans, G. A. Maree, J. Moolman, and R. J. Scholes. 2008. Designing protected areas to

674 conserve riverine biodiversity: lessons from a hypothetical redesign of the Kruger National
675 Park. *Biological Conservation* 141:100-117.

676

677 Saunders, D. L., J. J. Meeuwig, and A. C. J. Vincent. 2002. Freshwater protected areas: strategies
678 for conservation. *Conservation Biology* 16:30-41.

679

680 Sigler, J., and D. Zaroban. 2018. *Fishes of Idaho: a natural history survey*. Caxton Press.

681

682 Smith, G. R. 1978. Biogeography of Intermountain fishes. *Great Basin Naturalist Memoirs*
683 2(3):17-42.

684

685 Thurow, R. F., D. C. Lee, and B. E. Rieman. 1997. Distribution and status of seven native
686 salmonids in the interior Columbia River basin and portions of the Klamath River and Great
687 basins. *North American Journal of Fisheries Management* 17:1094–1110.

688

689 Thurow, R. F., D. C. Lee, and B. E. Rieman. 2000. Status and distribution of Chinook salmon and
690 steelhead in the interior Columbia River basin and portions of the Klamath River basin.
691 Pages 133–160 *in* Sustainable fisheries management: Pacific salmon. Knudsen, E., C.
692 Steward, D. MacDonald, J. Williams, and D. Reiser, editors. CRC Press, Boca Raton, Florida.

693

694 UDWR (Utah Division of Wildlife Resources). 2000. Cutthroat trout management: a position
695 paper. Genetic considerations associated with cutthroat trout management. Publication
696 Number 00-26, Developed by: Colorado Division of Wildlife; Idaho Department of Fish and
697 Game; Montana Fish, Wildlife and Parks; Nevada Division of Wildlife; New Mexico Game
698 and Fish; Utah Division of Wildlife Resources; Wyoming Game and Fish Department, Utah
699 Division of Wildlife Resources, Salt Lake City, Utah.

700

701 USGS. 2004. Land Ownership in Western North America, 180 m. Sage-grouse rangewide
702 conservation assessment, Snake River Field Station, U.S. Geological Survey, Boise, Idaho.

703

704 USGS (U.S. Geological Survey). 2011. Protected areas database of the United States (PAD-US)
705 Version 1.2. U.S. Geological Survey, Gap Analysis Program (GAP), Moscow, Idaho.

706

707 Wallace, R. L. and D. W. Zaroban. 2013. *Native fishes of Idaho*. American Fisheries Society,
708 Bethesda, Maryland.

709

710 Walrath, J. D., D. C. Dauwalter, and D. Reinke. 2016. Influence of stream condition on habitat
711 diversity and fish assemblages in an impaired upper Snake River Basin watershed.
712 *Transactions of the American Fisheries Society* 145:821-834.

713

714 Wenger, S. J., D. J. Isaak, C. H. Luce, H. M. Neville, K. D. Fausch, J. B. Dunham, D. C. Dauwalter,
715 M. K. Young, M. M. Elsner, B. E. Rieman, A. F. Hamlet, and J. E. Williams. 2011. Flow regime,
716 temperature, and biotic interactions drive differential declines of trout species under
717 climate change. *Proceedings of the National Academy of Sciences* 108:14175-14180.

718
719 Williams, J. E., A. L. Haak, N. G. Gillespie, and W. T. Colyer. 2007. The Conservation Success
720 Index: synthesizing and communicating salmonid condition and management needs.
721 Fisheries 32:477-492.
722
723 Williams, J. E., R. N. Williams, R. F. Thurow, L. Elwell, D. P. Phillipp, F. A. Harris, J. L. Kershner, P.
724 J. Martinez, D. Miller, G. H. Reeves, C. A. Frissell, and J. R. Sedell. 2011. Native Fish
725 Conservation Areas: a vision for large-scale conservation of native fish communities.
726 Fisheries 36:267-277.
727
728 Williams, J. E. 2019. Multispecies conservation: bringing efficiency to the science of native fish
729 conservation. Pages XXX - XXX in D. C. Dauwalter, T. W. Birdsong, and G. P. Garrett, editors.
730 Multispecies and watershed approaches to freshwater fish conservation. American Fisheries
731 Society, Bethesda, Maryland.
732
733 Zoellick, B. W., D. B. Allen, and B. J. Flatter. 2005. A long-term comparison of redband trout
734 distribution, density, and size structure in southwestern Idaho. North American Journal of
735 Fisheries Management 25:1179-1190.
736
737

738 Table 1. Fishes and other taxa native to the Upper Snake River basin above Hells Canyon and their general distribution (upper or lower basin relative to
 739 Shoshone Falls), Species of Greatest Conservation Need (SCGN) status, number of occurrences (and total sites) in survey data, and species weight used to rank
 740 subwatersheds.

Taxonomic group	Common name (species abbreviations)	Scientific name	Distribution in basin	SCGN	Occurrences (Total sites)	Analysis weight
Fishes	Pacific Lamprey (PLY)	<i>Entosphenus tridentate</i> ^b	Lower	ID, OR	Extirpated	
	White Sturgeon (WST)	<i>Acipenser transmontanus</i>	Lower	ID, OR		
	Chiselmouth (CSM)	<i>Acrocheilus alutaceus</i>	Lower		88 (1452)	0.945
	Utah Chub (UTC)	<i>Gila atraria</i>	Upper		23 (1280)	0.961
	Northern Leatherside Chub (NLC)	<i>Lepidomeda copei</i>	Upper	ID, WY	39 (1210)	0.977
	Peamouth (PMT)	<i>Mylocheilus caurinus</i>	Lower		No data	--
	Northern Pikeminnow (PMT)	<i>Ptychocheilus oregonensis</i>	Lower		123 (1448)	0.250
	Redside Shiner (RSS)	<i>Richardsonius balteatus</i>	Both		462 (3045)	0.371
	Longnose Dace (LND)	<i>Rhinichthys cataractae</i>	Both		298 (3002)	0.676
	Speckled Dace (SPD)	<i>Rhinichthys osculus</i>	Both		640 (3047)	0.001
	Kendall Warm Springs Dace (KWD)	<i>Rhinichthys osculus thermalis</i>	Upper	WY	No data	--
	Leopard Dace (LPD)	<i>Rhinichthys falcatus</i>	Lower	ID ^a	10 (660)	1.000
	Umatilla Dace (UMD)	<i>Rhinichthys umatilla</i>	Lower	ID ^a	No data	--
	Utah Sucker (UTS)	<i>Catostomus ardens</i>	Both		55 (1310)	0.938
	Bridgelip Sucker (BLS)	<i>Catostomus columbianus</i>	Lower		290 (1609)	0.250
	Wood River Bridgelip Sucker (WBLS)	<i>Catostomus columbianus hubbsi</i>	Lower		No data	--
	Bluehead Sucker (BHS)	<i>Catostomus discobolus</i>	Upper	ID ^a , UT, WY	46 (1296)	0.953
	Largescale Sucker (LSS)	<i>Catostomus macrocheilus</i>	Lower		101 (1609)	0.250
	Mountain Sucker (MTS)	<i>Catostomus platyrhynchus</i>	Upper		118 (2736)	0.820
	Snake River Sucker (SRS)	<i>Chasmistes muriei</i>	Upper		Extirpated	
	Yellowstone Cutthroat Trout (YCT)	<i>Oncorhynchus clarkii bouvieri</i>	Upper	ID, NV, WY		0.991
	Fine spotted form		Upper	ID, WY		0.736
	Redband (Rainbow) Trout (RBT)	<i>Oncorhynchus mykiss gairdneri</i>	Lower	ID, OR		0.923
	Summer Steelhead		Lower	ID, OR		Extirpated
	Bull Trout (BLT)	<i>Salvelinus confluentus</i>	Lower	ID, NV, OR		1.025
	Chinook Salmon (CHS)	<i>Oncorhynchus tshawytscha</i>	Lower	ID, NV, OR		Extirpated
	Sockeye Salmon (SES)	<i>Oncorhynchus nerka</i>	Lower	ID, OR		Extirpated
	Coho Salmon (COS)	<i>Oncorhynchus kisutch</i>	Lower	ID, OR		Extirpated
	Mountain Whitefish (MWF)	<i>Prosopium williamsoni</i>	Both	NV, WY	100 (1354)	0.558
	Mottled Sculpin (MSC)	<i>Cottus bairdii</i>	Both		263 (2628)	0.461
	Paiute Sculpin (PSC)	<i>Cottus beldingii</i>	Both		237 (2456)	0.340
	Shorthead Sculpin (SSC)	<i>Cottus confusus</i>	Lower		239 (1609)	0.719
	Shoshone Sculpin (ShSC)	<i>Cottus greenei</i>	Lower	ID	No data	--
Wood River Sculpin (WSC)	<i>Cottus leiopomus</i>	Lower	ID	44 (154)	0.922	
Crayfishes	Snake River pilose crayfish	<i>Pacifastacus connectens</i>	Both	ID, WY		

Shrimps	raptor fairy shrimp	<i>Branchinecta raptor</i>	Lower	ID
Snails	Banbury Springs lanx	<i>Lanx</i> sp.	Lower	ID
	pondsnail species group	<i>Stagnicola</i> sp.	Both	ID
	Rocky Mountain dusky snail	<i>Colligyrus greggi</i>	Upper	ID
	Bruneau Hot Spring snail	<i>Pyrgulopsis bruneauensis</i>	Lower	ID
	Bliss Rapids snail	<i>Taylorconcha serpenticola</i>	Both	ID
Mussels	California floater	<i>Anodonta californiensis</i>	Both	ID, NV, OR
	western ridged mussel	<i>Gonidea angulata</i>	Both	ID, OR
	western pearlshell	<i>Margaritifera falcata</i>	Both	ID
Salamander	western tiger salamander	<i>Ambystoma mavortium</i>	Upper	WY
Frogs	northern leopard frog	<i>Lithobates pipiens</i>	Both	ID, NV, WY
	Columbia spotted frog	<i>Rana luteiventris</i>	Lower	ID, NV, OR
Toads	Great Basin spadefoot	<i>Spea intermontana</i>	Both	NV, WY
	western toad	<i>Anaxyrus boreas</i>	Both	ID, NV, OR, WY
	Woodhouse toad	<i>Anaxyrus woodhouseii</i>	Lower	ID

^aSpecies listed as SGCN but status not assessed and available at time of analysis.

741
742
743
744

745 Table 2. Characteristics of potential NFCAs in the Upper Snake River basin. Species of Greatest Conservation Need shown in bold.

Watershed (subbasin)	Mean Rank (0 – 100)	Watershed area (km ²)	Native fishes	Protected (%) (stream corridor)	Public land (%)	Habitat Integrity ^b (5 – 25)	Future Security ^c (5 – 25)
Above Shoshone Falls							
Pacific Cr. (S. Fk. Snake)	98.8	431	YCT (both), NLC	95.4 (92.3)	97.3	25	21
Upper S. Fk. Snake R.	98.7	2,086	YCT (both), UTS, LND, SPD, RSS, MSC, PSC, UTC	99.9 (99.8)	92.2	24	21
Fall R. (Henry's Fork)	97.6	894	YCT (both), MTS, LND, SPD, RSS, MSC, PSC	93.4 (88.8)	97.9	23	16
Buffalo Fork (S. Fk. Snake)	95.8	959	YCT (both)	75.4 (80.2)	97.6	23	20
Goose Creek	95.3	1,842	YCT (lg.), BHS , BLS, MTS, UTS, LND, SPD, RSS, NLC , MSC, PSC, SSC, UTC	0.2 (0.1)	85.6	17	16
Conant Cr. (Henry's Fork)	95.0	312	YCT (both) BHS , MTS, LND, SPD, RSS, MSC, PSC	20.5 (8.8)	51.9	12	14
Cottonwood Cr. (S. Fk. Snake)	94.3	189	YCT (both), MWF	100 (100)	89.6	23	20
Spread Cr. (S. Fk. Snake)	90.6	344	YCT (both), PSC	15.3 (23.7)	90.7	19	19
Lower S. Fk. Snake R.	90.6	1,396	YCT (both), BHS , LSS, MTS, UTS, LND, SPD, RSS, MSC, PSC, MWF	0.7 (1.7)	74.7	16	13
McCoy Cr. (S. Fk. Snake)	89.5	282	YCT (both), MTS, UTS, LND, SPD, RSS, MSC, PSC	0.0 (0.0)	99.2	21	14
Ditch Cr. (S. Fk. Snake)	89.2	160	YCT (both), BHS	41.6 (10.6)	74.4	17	16
Gros Ventre R. (S. Fk. Snake)	88.7	1,617	YCT (both), PSC, MWF	36.4 (26.8)	97.6	20	18
Hoback R. (S. Fk. Snake)	87.9	1,469	YCT (fine), MTS, LND, MSC, MWF	19.3 (14.4)	94.1	21	14
Salt R. (S. Fk. Snake)	87.6	2,309	YCT (both), BHS , MTS, UTS, LND, SPD, RSS, NLC , MSC, PSC, MWF	0.7 (0.0)	71.3	15	15
Upper Raft R.	87.3	411	YCT (lg.), MTS, LND, SPD, RSS, MSC, PSC	4.9 (0.0)	30.2	18	16
Bear Cr. (S. Fk. Snake)	87.0	219	YCT (both), SPD, RSS, MSC, PSC	0.0 (0.0)	98.8	22	13
Upper Blackfoot R.	86.0	1,458	YCT (lg.), MTS, UTS, LND, SPD, RSS, MSC, PSC, UTC	0.4 (0.1)	56.0	16	15
Greys R. (S. Fk. Snake)	85.2	1,178	YCT (both), MTS, LND, MSC, PSC, MWF	0.3 (0.1)	99.7	19	16
Little Lost R.	84.3	1,528	BLT , MSC, SSC	10.7 (4.0)	94.7	17	21
Lower Blackfoot R.	83.3	1,832	YCT (lg.), MTS, UTS, LND, SPD, RSS, MSC, PSC, UTC	0.1 (0.4)	21.6	13	15
Big Elk Cr. (S. Fk. Snake)	81.6	160	YCT (both), SPD, MSC, PSC	0.0 (0.0)	97.6	24	14
Portneuf R.	80.0	4,344	YCT (lg.), BHS , MTS, UTS, LND, SPD, RSS, MSC, PSC, UTC	2.5 (0.1)	27.8	12	17
Willow Cr.	79.6	1,676	YCT (lg.), MTS, UTS, LND, SPD, RSS, MSC, PSC	19.0 (1.7)	36.9	15	17
Bitch Cr. (Henry's Fork)	78.7	245	YCT (lg.), MSC, PSC, MWF	77.2 (50.0)	78.6	22	14
Canyon Cr. (Henry's Fork)	77.4	215	YCT (lg.), LND, SPD, PSC, sculpin spp.	0.0 (0.0)	57.1	13	13
Cassia Cr. (Raft)	74.8	367	YCT (lg.), UTS, LND, RSS, MSC, PSC, UTC	0.4 (0.0)	74.8	15	17
Below Shoshone Falls							
Indian Cr. (Hells Canyon)	99.9	104	RBT, BLT	0.1 (0.0)	90.1	14	11
Little Jacks Cr. (Bruneau)	98.3	267	RBT , SPD, MSC, SSC	43.5 (59.2)	98.4	18	15
Brownlee Cr. (Hells Cyn.)	97.7	162	RBT , BLS, SSC, MWF	0.0 (0.0)	80.1	16	15
Squaw Cr. (Payette R.)	93.4	880	RBT , BLT , BLS, LSS, LND, SPD, RSS, NPM, MSC, SSC,	0.0 (0.0)	52.8	16	15
Lower N. Fk. Payette R.	93.1	810	RBT , LSS, SPD, RSS, NPM, MSC, SSC	0.0 (0.0)	50.3	15	12
Middle N. Fk. Payette R.	92.7	1,166	RBT , BLT , BLS, LSS, SPD, RSS, NPM, MSC	1.0 (0.4)	40.4	13	13
Middle Fk. Payette R.	92.3	878	RBT , BLT , CSM, LND, SPD, RSS, MSC, SSC, MWF	0.2 (0.3)	93.4	18	15
South Fk. Payette R.	90.0	1,918	RBT , BLT , LSS, CSM, LND, NPM, SSC	13.5 (10.7)	97.3	20	14
Upper Malheur R.	87.2	900	RBT , BLT , BLS, LSS, LND, SPD, RSS, NPM, MSC, MWF	18.8 (26.4)	76.3	15	18

South Fk. Boise R.	86.2	2,527	RBT, BLT , BLS, LSS, CSM, MTS, LND, SPD, RSS, NPM, MSC, SSC, MWF	0.0 (0.0)	88.4	17	16
Willow Cr. (Bruneau)	82.8	56	RBT	79.8 (80.5)	95.2	17	15
North/Middle Fk. Boise R.	79.7	3,156	RBT, BLT , BLS, LSS, LND, RSS, MSC, SSC, MWF	12.5 (12.3)	93.5	18	16
Jarbidge R. (Bruneau)	78.5	886	RBT, BLT , BLS, LSS, CSM, MTS, LND, SPD, RSS, NPM, MSC, SSC, MWF	47.7 (37.2)	89.6	18	16
Upper Little Weiser R.	78.4	205	RBT, BLT , LND, MSC, SSC	0.0 (0.0)	87.2	15	14
U. Deadwood R. (Payette)	77.5	283	RBT, BLT , SSC, MWF	0.0 (0.0)	95.5	21	15
Harrington Cr. (Owyhee)	69.2	151	RBT , SPD, PSC	0.0 (0.0)	61.0	16	16
Big Jacks (Bruneau)	65.3	632	RBT , BLS, LND, SPD, RSS, NPM	33.0 (66.1)	97.3	18	15
Cottonwood Cr. (Salmon Falls)	61.2	133	RBT , BLS, SPD, RSS, PSC	46.7 (42.5)	96.7	16	18

746 ^a See Table 1 for species abbreviations. Native fish occurrence based on project database and not expert knowledge.

747 ^b Habitat integrity from Trout Unlimited's (TU) Conservation Success Index. Subwatershed scores range from 5 (poor) to 25 (good). Habitat integrity based in indicators of
748 riparian condition, watershed connectivity, watershed condition, water quality, and flow regime.

749 ^c Future Security from TU's Conservation Success Index. Subwatershed scores range from 5 (not secure) to 25 (secure). Future security based in indicators of land conversion,
750 resource extraction, energy development, climate change, and introduced species.
751

752 Figure Captions

753

754 Figure 1. Conservation populations and current distributions of Bull Trout, Yellowstone
755 Cutthroat Trout (A), Redband Trout, and fine-spotted form of Yellowstone Cutthroat Trout (B)
756 in the Upper Snake River basin.

757

758 Figure 2. Conceptual model showing data integration for subwatershed ranking analysis.

759

760 Figure 3. Subwatershed ranks in the Upper Snake River basin based on native trout
761 distributions and abundance, native non-game species occurrence probability, drainage
762 network connectivity, and land protection status.

763

764 Figure 4. Potential NFCAs in the Upper Snake River Basin with the presence of non-game
765 Species of Greatest Conservation Need (A) and pie charts illustrating the land status of
766 perennial stream corridors (B). Details of each watershed are found in Table 2.

767


768 Figure 5. Spatial distribution of Habitat Integrity (A) and Future Security (B) scores for
769 subwatersheds (HUC 12: n = 2079) in the Upper Snake River basin above Hells Canyon. Habitat
770 Integrity and Future Security indicators were based on Trout Unlimited's Conservation Success
771 Index (Williams et al. 2007).

772


773 Figure 6. Land ownership and native trout distributions in the Jarbidge River (A), Goose Creek
774 (B), and Upper Blackfoot River (C) watersheds.

775


A)



B)



GIS data pre-processing




ZONATION input files

Species weights

Spp	wt.
1	0.4
2	1.0
:	:

Subwatershed connectivity

Up	down
1	3
2	3
:	:



ZONATION automated pre-processing

Cost


Iterative ZONATION ranking

Additive Benefit Function


$$\delta_i = \frac{1}{c_i} w_j \sum_j \Delta V_j$$

Output


Subwatershed rankings (0 – 100)




A)




B)





A)



B)





Legend

Barriers

- Complete
- Partial
- Unknown

- Redband trout
- Bull trout
- Redband and bull trout
- Y. cutthroat trout

Land ownership

- Forest Service
- Bureau of Land Management
- National Park Service
- Fish and Wildlife Service
- State
- Other